

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LAGUNA 430

FUNGICIDE

ACTIVE CONSTITUENT: 430 g/L TEBUCONAZOLE

GROUP	3	FUNGICIDE
-------	----------	-----------

For the control of leaf spot and leaf speckle on bananas, rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum, and ryegrass and fescue seed crops as specified in the directions for use table.

CONTENTS: 5, 10, 20, 110, 1000 LITRES

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

APVMA Approval No: 59420/0105



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

Laguna 430 Fungicide

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with the eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA 131126)

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

Batch No:

Date of Manufacture:

This product is NOT a Dangerous Good under the Australian Code for the Transport of Dangerous Goods by road or rail.
For specialist advice in emergency only, call 1800 033 111 all hours, Australia wide



G.H.S. STATEMENT

HAZARDS H361: Suspected of damaging fertility or the unborn child. H410: Very toxic to aquatic life with long lasting effects.

PREVENTION P270: Do not eat, drink or smoke when using this product. P280: Wear protective gloves, protective clothing and eye or face protection.

RESPONSE P301+P330+P331: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.

Continue rinsing. P370+P378: In case of fire, note the following. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used. Try to contain spills, minimise spillage entering drains or water courses.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LAGUNA 430

FUNGICIDE

ACTIVE CONSTITUENT: 430 g/L TEBUCONAZOLE

GROUP	3	FUNGICIDE
-------	----------	-----------

For the control of leaf spot and leaf speckle on bananas, rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum, and ryegrass and fescue seed crops as specified in the directions for use table.

THIS LEAFLET IS PART OF THE LABEL

APVMA Approval No: 59420/0105



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

Directions for use:

Crop	Disease	Application	Rate	WHP	Critical Comments
Bananas (QLD NSW WA NT ONLY)	Leaf spot (yellow sigatoka) Leaf speckle Black sigatoka	Ground Application Apply by misting machine or airblast sprayer in a convenient volume of water Aerial application Apply at least 20 L of spray mixture per hectare	230 mL/ha Add 3-6 L water miscible oil/ha	1 day	Maintain good deleafing practices to reduce disease inoculum. Very old leaves and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of TEBUCONZOLE. Tropical areas (eg North Qld, NT, Ord River): Apply a regular schedule of protectant sprays. When conditions favour disease, apply a minimum of 2 and a maximum of 3 consecutive LAGUNA 430 FUNGICIDE sprays at 14 day intervals. Do not apply more than 6 LAGUNA 430 FUNGICIDE sprays in any 12 month period. Do not apply any LAGUNA 430 FUNGICIDE sprays in the months of July, August and September. Sub-tropical areas (eg South Qld, NSW): Commence spraying with LAGUNA 430 FUNGICIDE at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a minimum of 2 consecutive LAGUNA 430 FUNGICIDE sprays. Do not apply more than 5 LAGUNA 430 FUNGICIDE sprays in any 12 month period. LAGUNA 430 FUNGICIDE is approved for use in banana plantations interplanted with avocados.
Peanuts (South QLD NSW ONLY)	Early leaf spot. Late Leaf spot Rust Net blotch	Low disease Ground Application Apply in at least 100 L of water/ha. Aerial application Apply at least 30 L of spray mixture per hectare.	175 mL/ha + Argridex at 1 L/ha	H3 weeks G3 weeks	Regularly check high risk areas in the crop for disease (eg lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found then either: -spray immediately after the last disease conducive weather (eg rain or heavy dews) OR – spray before the next disease conducive weather. Repeat after 14 days if conditions remain favourable to disease development. If not, repeat before or just after the next disease conducive weather. For resistance strategy, see General Instructions.
		High disease OR wet weather	290 mL/ha + Agridex 1 L/ha		
		High disease AND wet weather	440 mL/ha + Agridex 1 L/ha		

Crop	Disease	Application	Rate	WHP	Critical Comments	
Peanuts (North QLD WA NT ONLY)	Early leaf spot, Late leaf spot, Rust	Low disease	230 mL/ha + Agridex 1 L/ha.	H3 weeks G3 weeks	Begin spraying at 3-4 weeks after planting. If band spraying apply half the appropriate rate as a 45cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14 day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10-12 days. For resistance strategy see General Instructions.	
		Moderate disease	290 mL/ha + Agridex 1 L/ha.			
		Severe disease	440 mL/ha + Agridex 1 L/ha			
	Net blotch	Low disease	290 mL/ha + Agridex 1 L/ha.			Apply at 14 day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.
	Moderate – severe disease	440 mL/ha + Agridex L/ha				
Green beans	Rust	Can be applied by aircraft or ground rig	350 mL/ha + Agridex 1 L/ha	H 3 days G3 days	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high.	
Peas	Powdery mildew	Ground application: Apply in at least 50 L of water/ha	145 mL/ha		Apply at flowering or at first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions.	
Wheat	Leaf rust, Stripe rust, Septoria nodorum blotch, Yellow leaf spot	Aerial application: Apply in at least 10 L of water/ha	145 or 290 mL/ha	H 5 weeks G 14 days	Use higher rate when longer disease control is required. Stripe rust: See spray timings under General Instructions. Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil eg D-C- Trate or equivalent) at 1%, may improve performance of LAGUNA 430 FUNGICIDE on wheat, oats and barley.	
	Septoria tritici blotch		290 mL/ha			
Oats	Crown rust		145 or 290 mL/ha			
Barley	Scald		145 mL/ha			Apply at late tillering to early jointing.
	Powdery mildew		145 or 290 mL/ha			Apply when 5% of the leaf area is infected. Use higher rate when longer disease control is required.
Wheat Oats	Stem rust					145 or 290mL/ha

Crop	Disease	Application	Rate	WHP	Critical Comments
Onions (TAS ONLY)	White root rot	Before sowing, apply Laguna 430 Fungicide onto lime super. Ensure good coverage of all lime super particles.	1.45 mL / 100 m of row mixed with 145- 218 g lime super/100 m of row	-	Apply LAGUNA 430 FUNGICIDE treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a band width of 2 cm. Ensure that the correct rate of LAGUNA 430 FUNGICIDE is used otherwise some delay in emergence and reduced stands of seedlings may occur.
Pawpaw	Black spot	Ensure thorough coverage of leaves and fruit.	290 ml/ha	3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of LAGUNA 430 FUNGICIDE. Apply LAGUNA 430 FUNGICIDE at 14 day intervals. Alternate LAGUNA 430 FUNGICIDE with sprays of a protectant fungicide (eg Dithane DF). Do not apply more than 6 sprays of LAGUNA 430 FUNGICIDE (or any DMI fungicide) on any block in any 12 month period.
Pyrethrum	<i>Sclerotinia sclerotior um</i>		350 mL/ha	-	Apply twice, in rotation with other control measures at 7-10 day intervals. Commence at 1 to 2% flowering. Use under directions of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf rust Stem rust	Apply in at least 100 L of water/ha	290 mL/ha		Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or reinfection may require a further application 3-4 weeks later. Ensure thorough coverage, and use higher water volumes in dense or advanced crops.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

BANANAS, AVOCADOS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

BEANS, PEAS: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

BEANS, PEAS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 DAYS AFTER APPLICATION.

CEREALS: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

CEREALS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

ONIONS: NOT REQUIRED WHEN USED AS DIRECTED.

PAPAW: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

PEANUTS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

PEANUTS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

H = harvest, G = grazing.

GENERAL INSTRUCTIONS

GROUP	C	FUNGICIDE
-------	---	-----------

FUNGICIDE RESISTANCE WARNING

Laguna 430 Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist though normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resisting fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

RESISTANCE MANAGEMENT RECOMMENDATION - PEANUTS

Apply no more than 3 consecutive sprays of DMI fungicide (eg TEBUCONAZOLE) before switching to a non-DMI fungicide. Apply no more than 5 DMI sprays per season.

SPECIAL WARNING - BANANAS

There are certain conditions when the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise risk of fruit marking. **Never include adjuvants** other than water miscible oils with LAGUNA 430 FUNGICIDE sprays. The inclusion of wetting agents is known to cause phototoxicity to young fruit.

FOLIAR DISEASES ON CEREAL CROPS

Do not apply to cereal crops more than once in a season. Treatment will give approximately three weeks disease suppression. Economic responses may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7-10 days after application.

Yield potential: crops with potential yield under 2 t/ha are unlikely to give economic responses to a fungicide spray except under conditions of very severe disease. Economic responses are most likely with crops with potential yield of over 3 t/ha.

SPRAY TIMINGS FOR STRIPE RUST CONTROL

Obtain advisory literature from Department of Agriculture for classification of resistant and susceptible varieties. In South Australia consult Plant Protection Note PPN 21. In other states the following spray program is suggested:

Seedling infections: When approximately 20 out of 100 leaves show first signs of infection during tillering to jointing, apply a spray within one week.

Adult infections – susceptible varieties: When approximately 10 out of 100 leaves show first signs of infection, apply within one week. Do not delay.

Adult infections – moderately susceptible varieties: When approximately 15 to 20 leaves out of 100 leaves show first signs of infection, apply a spray within one week. Do not delay.

Adult infections – moderately resistant and resistant varieties: Monitor carefully. If rust appears and spreads, spray within one week.

MIXING

Prior to pouring, shake container vigorously, then add the required amount of Laguna 430 Fungicide to water in the spray vat while stirring or with agitators in motion. Add the required amount of Agridex (peanuts, beans) or water miscible oil (bananas) and mix thoroughly.

APPLICATION

Aircraft should fly as low as possible under the prevailing conditions to minimise drift.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, ponds, waterways or drains with the product or used container.

The spray drift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practices Manual.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with the eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE AUSTRALIA 131126)

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.