

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PYRESTA® XTREME®

LV HERBICIDE

Active Constituents: 2.1 g/L PYRAFLUFEN-ETHYL 600 g/L 2, 4-D (present as the 2-ethylhexyl ester)



A low volatile herbicide for improvement in the brownout of a range of broadleaf and grass weeds, and improved control of marshmallow, long storksbill (*Erodium*) and wild radish, when used in tank mixtures with Raze®, Firebolt® or other glyphosate based herbicides as per the Directions for Use Table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

Contents: 20L - 110L

APVMA Approval No: 69952/62158



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

PYRESTA XTREME LV HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots in compliance with relevant local state or territory government legislation. Do not burn empty containers or product.

Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with the eyes. Avoid inhaling. When opening the container, preparing spray and using prepared spray, wear cotton overalls, over normal clothing buttoned to the neck and wrist, a washable hat or elbow length chemical resistant gloves. In addition, when mixing and loading, wear face shield or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each days use wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au.

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

Batch No:

Date of Manufacture:



This product is NOT a Dangerous Good under the Australian Code for the transport of Dangerous Goods by road and rail.

**For specialist advice in emergency only, call
1800 033 111 all hours, Australia wide**

G.H.S. STATEMENTS

HAZARDS H335: May cause respiratory irritation.

PREVENTION P262: Do not get in eyes, on skin, or on clothing.

P264: Wash contacted areas thoroughly after handling. P270: Do not eat, drink or smoke when using this product. P271: Use only outdoors or in a well ventilated area. P272: Contaminated work clothing should not be allowed out of the workplace.

RESPONSE P301+P330+P331: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. P302+P352: IF ON SKIN: Wash with plenty of soap and water. P304+P340: IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PYRESTA® XTREME®

LV HERBICIDE

Active Constituents: 2.1 g/L PYRAFLUFEN-ETHYL 600 g/L 2, 4-D (present as the 2-ethylhexyl ester)



A low volatile herbicide for improvement in the brownout of a range of broadleaf and grass weeds, and improved control of marshmallow, long storksbill (*Erodium*) and wild radish, when used in tank mixtures with Raze®, Firebolt® or other glyphosate based herbicides as per the Directions for Use Table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.

THIS LEAFLET IS PART OF THE LABEL

APVMA Approval No: 69952/62158



SIPCAM PACIFIC AUSTRALIA PTY LIMITED
Level 1, 191 Malop Street Geelong VIC 3220

® Registered Trademark of Sipcam Pacific Australia Pty Ltd

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT use open mixing/loading unless application is by a closed cab vehicle.

DO NOT apply to cotton by ground boom equipment.

DO NOT apply by low pressure hand wand equipment.

DO NOT apply using spraying equipment carried on the back of the user.

DO NOT apply using aerial spraying equipment.

AVOID spraying if rain is likely within 6 hours or if strong winds prevail.

DO NOT apply if weeds are stressed due to dry or excessively moist conditions.

DO NOT use when breeze is blowing towards nearby susceptible plants.

Situation & Crop	State	Weeds Controlled	Rate/ha	Critical Comments
Prior to sowing broadacre crops or starting a fallow (Winter weeds)	All States	Amsinckia (<i>Amsinckia</i> spp) Brome grass (<i>Bromus</i> spp) Capeweed (<i>Arctotheca calendula</i>) Chickweed (<i>Stellaria media</i>) Deadnettle (<i>Lamium amplexicaule</i>) Long storksbill (<i>Erodium botrys</i>) Medic (<i>Medicago</i> spp) Paterson's curse (<i>Echium plantagineum</i>) Turnip weed (<i>Rapistrum rugosum</i>) Volunteer barley Wild radish (<i>Raphanus raphanistrum</i>) Wild turnip (<i>Brassica tournefortii</i>)	250 – 500 mL plus recommended rate of Raze or other glyphosate herbicide	For improvement in the brownout of a range of broadleaf and grass weeds, and improved control of marshmallow, long storksbill (<i>Erodium</i>) and wild radish, when used in tank mixtures with Raze®, Firebolt® or other glyphosate based herbicides. Refer to the Raze, Firebolt or other glyphosate product label and select a rate suitable for the conditions and weed size in each situation. Apply when weeds are actively growing and at the 2-6 leaf growth stage. The higher rate of Pyresta Xtreme will provide the maximum speed of brownout and control. Visible symptoms of brownout may take from 2-7 days to develop.
		Above weeds plus: Annual ryegrass (<i>Lolium rigidum</i>) Barley grass (<i>Hordeum leporinum</i>) Curled dock (<i>Rumex crispus</i>) Indian hedge mustard (<i>Sisymbrium orientale</i>) Silver grass (<i>Vulpia</i> spp) Sowthistle (<i>Sonchus oleraceus</i>) Subterranean clover (<i>Trifolium subterraneum</i>) Volunteer wheat Wild oats (<i>Avena</i> spp)	500 mL plus recommended rate of Raze or other glyphosate herbicide	
		Above weeds plus: Marshmallow (<i>Malva</i> spp)	500 - 900 mL plus recommended rate of Raze or other glyphosate herbicide	
Prior to sowing broadacre crops or starting a fallow (Summer weeds)	Southern NSW, Vic, SA, WA only	Bathurst burr (<i>Xanthium spinosum</i>), Caltrop (<i>Tribulus terrestris</i>), Pigweed (<i>Portulaca oleracea</i>) Prickly paddy melon (<i>Cucumis myriocarpus</i>) Wild (Afghan) melon (<i>Citrullus lanatus</i>)	500 - 900 mL plus recommended rate of Raze or other glyphosate herbicide	Application of Pyresta Xtreme to old, hardened or stressed plants and in harsh environmental conditions such as high temperatures and low soil moisture may not provide adequate control. Where Barnyard grass is present high rates of Raze Herbicide may be required to avoid antagonistic effects.
		Volunteer Chickpea (<i>Cicer arietinum</i>)	900 mL plus recommended rate of Raze or other glyphosate herbicide	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIOD (WHP):

DO NOT GRAZE TREATED WEEDS FOR 7 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions for Pyresta Xtreme and for the glyphosate product to be added to Pyresta Xtreme.

The addition of Pyresta Xtreme to Raze, Firebolt or other glyphosate products may increase the rate of brownout and final control, when compared to the use of the glyphosate products alone. Refer to the product label for Raze, Firebolt or other glyphosate product and select a rate suitable for the conditions and weed size in each situation.

MIXING AND APPLICATION

Apply by ground application only. Half fill the spray tank with good quality water and begin agitation. While agitating, add the required volume of Pyresta Xtreme then the required volume of Raze, Firebolt or other glyphosate product, followed by the remaining water. Prepare only enough spray solution for immediate use, and do not stand overnight. If tank mixtures are prepared maintain good agitation and use promptly.

Spray equipment should be properly calibrated to ensure correct and uniform coverage. Use sufficient good quality water and configure the boom sprayer to optimise coverage. Apply in 60 to 150 L water per hectare.

DO NOT apply with smaller than Coarse to Very Coarse spray droplets according to the ASAE S572 definition for standard nozzles.

EQUIPMENT CLEANING

Do not allow the spray solution to dry in the application equipment. After application and before using the sprayer equipment for any other applications, the sprayer must be thoroughly cleaned. Applicators must ensure proper equipment clean-out for any other products mixed with Pyresta Xtreme as provided on the other product label(s). Immediately following application, clean all equipment thoroughly with detergent or a spray tank cleaner and water as described below. Should residues of Pyresta Xtreme remain in inadequately cleaned equipment, they may be released in subsequent applications and cause injury to crops.

1. Drain sprayer tank, hoses and spray boom and thoroughly rinse with clean water the inside of the spray tank, sprayer hoses, boom, and nozzles to remove any sediment or residues. Drain in an area well clear from watercourses, desirable vegetation and tree roots, that will also not drain towards watercourses, desirable vegetation and tree roots.
2. Fill the tank with clean water, add an alkaline detergent or boom cleaner (follow manufacturer's directions for use). Fill tank to capacity and operate the sprayer with agitation for 15 minutes to flush hoses, boom and nozzles.
3. Drain the sprayer tank, lines, and booms. Rinse the tank with clean water and flush through the hoses, boom and nozzles. Remove and clean spray nozzles, tips and screens.

COMPATIBILITY

Pyresta Xtreme is compatible with Raze, Firebolt and other formulations of glyphosate.

RESISTANT WEEDS WARNING

GROUP	G	I	HERBICIDE
-------	---	---	-----------

Pyresta Xtreme LV Herbicide is a member of both the aryl triazolinone and phenoxy group of herbicides. Its modes of action are through a process of membrane disruption, which is initiated by the inhibition of the enzyme protoporphyrinogen oxidase and through disruptors of plant cell growth. For weed resistance management, Pyresta Xtreme belongs to the Group G and Group I herbicides. Some naturally occurring weed biotypes resistant to Pyresta Xtreme and other Group G and I herbicides may exist through normal genetic variability in any weed population and increase if these herbicides are used repeatedly. These resistant weeds will not be controlled by Pyresta Xtreme or other Group G or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcarn Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Pyresta Xtreme or other Group G or Group I herbicides.

PLANTBACK AND ROTATION RECOMMENDATIONS

Plant back days for Pyresta Xtreme LV Herbicide

Crop	250 to 500 mL/ha	900 mL/ha
Balansa Clover	7	7
Barley%	1	1
Chickpeas#	7	14
Faba Beans	7	7
Field Peas	7	14
Lentils	7	7
Linseed	7	7
Lucerne	7	7
Lupins+	7	14
Medics	7	7
Narbon Beans	7	7
Navybean	10	10
Oats	3	3
Perennial Ryegrass	7	7
Persian Clover	7	7
Phalaris	7	7
Canola/Rapeseed#	14	21
Rice	7	7
Safflower#	7	14
Sorghum@	3	7
Soybean	14	14
Sub-Clover	7	7
Sunflower@	7	10
Triticale%	1	3
Vetch	7	7
Wheat%	1	3
White Clover	7	7

IMPORTANT:

WHEN APPLIED TO DRY SOILS AT LEAST 15 MM OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for wheat, barley and triticale.

In Queensland, planting of canola/rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15 mm.

@ In Central Queensland, the Plant Back Period for sorghum and sunflower is 1 day irrespective of rainfall.

+ In Western Australia, the Plant Back Period for lupins is 28 days.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (over normal clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds.

Drift Warning:

DO NOT use unless wind speed is more than 3 kilometers per hour and less than 15 kilometers per hour as measured at the application site. DO NOT apply with smaller than Coarse to Very Coarse spray droplets according to the ASAE S572 definition for standard nozzles. Direct spray contact or even slight drift may cause severe injury or destruction of any growing crop or other desirable plants including trees and native vegetation. DO NOT use when breeze is blowing towards nearby desirable plants. DO NOT apply under meteorological conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. Equipment settings which produce fine droplets (150 microns or less), "still air" and hot dry days all contribute to drift. This is a phenoxy herbicide that can cause severe damage to susceptible crops such as cotton, tobacco, tomatoes, flowers, vegetables, vines, fruit trees, legume crops and pastures, oil seed crops or other susceptible crops and trees (eg Kurrajongs, Belahs, Eucalypts).

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. Do not contaminate wetlands or waterways with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots in compliance with relevant local state or territory government legislation. Do not burn empty containers or product.

Refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with the eyes. Avoid inhaling. When opening the container, preparing spray and using prepared spray, wear cotton overalls, over normal clothing buttoned to the neck and wrist, a washable hat or elbow length chemical resistant gloves. In addition, when mixing and loading, wear face shield or goggles. If product in eyes, wash out immediately with water. Wash hands after use. After each days use wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au.

NOTICE TO BUYER

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.